

BOUNDS FOR THE COMBINATION OF TOADER MEAN AND THE ARITHMETIC MEAN IN TERMS OF THE CONTRAHARMONIC MEAN

WEI-DONG JIANG AND FENG QI

ABSTRACT. In the paper, the authors find the greatest value λ and the least value μ such that the double inequality

$$C(\lambda a + (1 - \lambda)b, \lambda b + (1 - \lambda)a) < \alpha A(a, b) + (1 - \alpha)T(a, b) \\ < C(\mu a + (1 - \mu)b, \mu b + (1 - \mu)a)$$

holds for all $\alpha \in (0, 1)$ and $a, b > 0$ with $a \neq b$, where

$$C(a, b) = \frac{a^2 + b^2}{a + b}, \quad A(a, b) = \frac{a + b}{2},$$

and

$$T(a, b) = \frac{2}{\pi} \int_0^{\pi/2} \sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta} \, d\theta$$

denote respectively the contraharmonic, arithmetic, and Toader means of two positive numbers a and b .

1. INTRODUCTION

For $p \in \mathbb{R}$ and $a, b > 0$, the contraharmonic mean $C(a, b)$, the p -th power mean $M_p(a, b)$, and Toader mean $T(a, b)$ are respectively defined by

$$C(a, b) = \frac{a^2 + b^2}{a + b}, \quad M_p(a, b) = \begin{cases} \left(\frac{a^p + b^p}{2} \right)^{1/p}, & p \neq 0, \\ \sqrt{ab}, & p = 0, \end{cases}$$

and

$$T(a, b) = \frac{2}{\pi} \int_0^{\pi/2} \sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta} \, d\theta = \begin{cases} \frac{2a}{\pi} \mathcal{E} \left(\sqrt{1 - \left(\frac{b}{a} \right)^2} \right), & a > b, \\ \frac{2b}{\pi} \mathcal{E} \left(\sqrt{1 - \left(\frac{b}{a} \right)^2} \right), & a < b, \\ a, & a = b, \end{cases} \quad (1.1)$$

where

$$\mathcal{E} = \mathcal{E}(r) = \int_0^{\pi/2} \sqrt{1 - r^2 \sin^2 \theta} \, d\theta$$

2010 *Mathematics Subject Classification.* 26E60, 33E05.

Key words and phrases. bound; contraharmonic mean; arithmetic mean; Toader mean; complete elliptic integrals.

This paper was typeset using $\mathcal{A}_{\mathcal{M}}\mathcal{S}\text{-}\mathcal{L}^{\text{A}}\mathcal{T}_{\text{E}}\mathcal{X}$.

for $r \in [0, 1]$ is the complete elliptic integral of the second kind. For more information on complete elliptic integrals, please see [11, 12, 13, 14] and plenty of references therein.

Recently, Toader mean has attracted attention of several researchers. In particular, many remarkable inequalities for $T(a, b)$ can be found in the literature [6, 7, 9, 10, 17]. It was conjectured in [16] that

$$M_{3/2}(a, b) < T(a, b) \quad (1.2)$$

for all $a, b > 0$ with $a \neq b$. This conjecture was proved in [3, 15] respectively. In [1], a best possible upper bound for Toader mean was presented by

$$T(a, b) < M_{\ln 2 / \ln(\pi/2)}(a, b) \quad (1.3)$$

for all $a, b > 0$ with $a \neq b$.

It is not difficult to verify that

$$C(a, b) > M_2(a, b) = \sqrt{\frac{a^2 + b^2}{2}} \quad (1.4)$$

for all $a, b > 0$ with $a \neq b$. From (1.2) to (1.4) one has

$$A(a, b) < T(a, b) < C(a, b)$$

for all $a, b > 0$ with $a \neq b$.

For positive numbers $a, b > 0$ with $a \neq b$, let

$$J(x) = C(xa + (1-x)b, xb + (1-x)a) \quad (1.5)$$

on $[\frac{1}{2}, 1]$. It is not difficult to verify that $J(x)$ is continuous and strictly increasing on $[\frac{1}{2}, 1]$. Note that $J(\frac{1}{2}) = A(a, b) < T(a, b)$ and $J(1) = C(a, b) > T(a, b)$.

In [8] it was proved that the double inequality

$$C(\alpha a + (1-\alpha)b, \alpha b + (1-\alpha)a) < T(a, b) < C(\beta a + (1-\beta)b, \beta b + (1-\beta)a) \quad (1.6)$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha \leq \frac{3}{4}$ and $\beta \geq \frac{1}{2} + \frac{\sqrt{4\pi - \pi^2}}{2\pi}$.

The main purpose of the paper is to find the greatest value λ and the least value μ such that the double inequality

$$\begin{aligned} C(\lambda a + (1-\lambda)b, \lambda b + (1-\lambda)a) &< \alpha A(a, b) + (1-\alpha)T(a, b) \\ &< C(\mu a + (1-\mu)b, \mu b + (1-\mu)a) \end{aligned} \quad (1.7)$$

holds for all $\alpha \in (0, 1)$ and $a, b > 0$ with $a \neq b$. As applications, we also present new bounds for the complete elliptic integral of the second kind.

2. PRELIMINARIES AND LEMMAS

In order to establish our main result, we need several formulas and lemmas below.

For $0 < r < 1$ and $r' = \sqrt{1-r^2}$, Legendre's complete elliptic integrals of the first and second kinds are defined in [4, 5] by

$$\begin{cases} \mathcal{K} = \mathcal{K}(r) = \int_0^{\pi/2} \frac{1}{(1-r^2 \sin^2 \theta)^{1/2}} d\theta, \\ \mathcal{K}' = \mathcal{K}'(r) = \mathcal{K}(r'), \\ \mathcal{K}(0) = \frac{\pi}{2}, \\ \mathcal{K}(1) = \infty \end{cases}$$

and

$$\begin{cases} \mathcal{E} = \mathcal{E}(r) = \int_0^{\pi/2} (1 - r^2 \sin^2 \theta)^{1/2} d\theta, \\ \mathcal{E}' = \mathcal{E}'(r) = \mathcal{E}(r'), \\ \mathcal{E}(0) = \frac{\pi}{2}, \\ \mathcal{E}(1) = 1 \end{cases}$$

respectively.

For $0 < r < 1$, the formulas

$$\begin{aligned} \frac{d\mathcal{K}}{dr} &= \frac{\mathcal{E} - (r')^2 \mathcal{K}}{r(r')^2}, \quad \frac{d\mathcal{E}}{dr} = \frac{\mathcal{E} - \mathcal{K}}{r}, \quad \frac{d(\mathcal{E} - (r')^2 \mathcal{K})}{dr} = r\mathcal{K}, \\ \frac{d(\mathcal{K} - \mathcal{E})}{dr} &= \frac{r\mathcal{E}}{(r')^2}, \quad \mathcal{E}\left(\frac{2\sqrt{r}}{1+r}\right) = \frac{2\mathcal{E} - (r')^2 \mathcal{K}}{1+r} \end{aligned}$$

were presented in [2, Appendix E, pp. 474–475].

Lemma 2.1 ([2, Theorem 3.21(1) and 3.43 Exercise 13(a)]). *The function $\frac{\mathcal{E} - (r')^2 \mathcal{K}}{r^2}$ is strictly increasing from $(0, 1)$ onto $(\frac{\pi}{4}, 1)$ and the function $2\mathcal{E} - (r')^2 \mathcal{K}$ is increasing from $(0, 1)$ onto $(\frac{\pi}{2}, 2)$.*

Lemma 2.2. *Let $u, \alpha \in (0, 1)$ and*

$$f_{u,\alpha}(r) = ur^2 - (1 - \alpha) \left\{ \frac{2}{\pi} [2\mathcal{E}(r) - (1 - r^2)\mathcal{K}(r)] - 1 \right\}. \quad (2.1)$$

Then $f_{u,\alpha} > 0$ for all $r \in (0, 1)$ if and only if $u \geq (1 - \alpha)(\frac{4}{\pi} - 1)$ and $f_{u,\alpha} < 0$ for all $r \in (0, 1)$ if and only if $u \leq \frac{1-\alpha}{4}$.

Proof. It is clear that

$$f_{u,\alpha}(0^+) = 0, \quad (2.2)$$

$$f_{u,\alpha}(1^-) = u - (1 - \alpha) \left(\frac{4}{\pi} - 1 \right), \quad (2.3)$$

$$f'_{u,\alpha}(r) = 2r[u - (1 - \alpha)g(r)], \quad (2.4)$$

where $g(r) = \frac{1}{\pi} \frac{\mathcal{E} - (r')^2 \mathcal{K}}{r^2}$.

When $u \geq \frac{1-\alpha}{\pi}$, from (2.4) and Lemma 2.1 and by the monotonicity of $g(r)$, it follows that $f_{u,\alpha}(r)$ is strictly increasing on $(0, 1)$. Therefore, $f_{u,\alpha}(r) > 0$ for all $r \in (0, 1)$.

When $u \leq \frac{1-\alpha}{4}$, from (2.4) and Lemma 2.1 and by the monotonicity of $g(r)$, we obtain that $f_{u,\alpha}(r)$ is strictly decreasing on $(0, 1)$. Therefore, $f_{u,\alpha}(r) < 0$ for all $r \in (0, 1)$.

When $\frac{1-\alpha}{4} < u \leq (1 - \alpha)(\frac{4}{\pi} - 1)$, from (2.3) and (2.4) and by the monotonicity of $g(r)$, we see that there exists $\lambda \in (0, 1)$ such that $f_{u,\alpha}(r)$ is strictly increasing in $(0, \lambda]$ and strictly decreasing in $[\lambda, 1)$ and

$$f_{u,\alpha}(1^-) \leq 0. \quad (2.5)$$

Therefore, making use of the equation (2.2), the inequality (2.5), and the piecewise monotonicity of $f_{u,\alpha}(r)$ lead to the conclusion that there exists $0 < \lambda < \eta < 1$ such that $f_{u,\alpha}(r) > 0$ for $r \in (0, \eta)$ and $f_{u,\alpha}(r) < 0$ for $r \in (\eta, 1)$.

When $(1 - \alpha)(\frac{4}{\pi} - 1) \leq u < \frac{1-\alpha}{\pi}$, by (2.3), it follows that

$$f_{u,\alpha}(1^-) \geq 0. \quad (2.6)$$

From (2.3) and (2.4) and by the monotonicity of $g(r)$, we see that there exists $\lambda \in (0, 1)$ such that $f_{u,\alpha}(r)$ is strictly increasing in $(0, \lambda]$ and strictly decreasing in $[\lambda, 1)$. Therefore, $f_{u,\alpha}(r) > 0$ for $r \in (0, 1)$ follows from (2.2) and (2.6) together with the piecewise monotonicity of $f_{u,\alpha}(r)$. \square

3. MAIN RESULTS

Now we are in a position to state and prove our main results.

Theorem 3.1. *If $\alpha \in (0, 1)$ and $\lambda, \mu \in (\frac{1}{2}, 1)$, then the double inequality*

$$\begin{aligned} C(\lambda a + (1 - \lambda)b, \lambda b + (1 - \lambda)a) &< \alpha A(a, b) + (1 - \alpha)T(a, b) \\ &< C(\mu a + (1 - \mu)b, \mu b + (1 - \mu)a) \end{aligned} \quad (3.1)$$

holds for all $a, b > 0$ with $a \neq b$ if and only if

$$\lambda \leq \frac{1}{2} + \frac{\sqrt{1-\alpha}}{4} \quad \text{and} \quad \mu \geq \frac{1}{2} \left[1 + \sqrt{(1-\alpha) \left(\frac{4}{\pi} - 1 \right)} \right].$$

Proof. Since $A(a, b)$, $T(a, b)$, and $C(a, b)$ are symmetric and homogeneous of degree one, without loss of generality, assume that $a > b$. Let $p \in (\frac{1}{2}, 1)$, $t = \frac{b}{a} \in (0, 1)$, and $r = \frac{1-t}{1+t}$. Then

$$\begin{aligned} &C(pa + (1-p)b, pb + (1-p)a) - \alpha A(a, b) - (1-\alpha)T(a, b) \\ &= a \frac{[p + (1-p)b/a]^2 + (pb/a + 1-p)^2}{1+b/a} - \alpha a \frac{1+b/a}{2} - (1-\alpha) \frac{2a}{\pi} \mathcal{E} \left(\sqrt{1 - \left(\frac{b}{a} \right)^2} \right) \\ &= a \left\{ \frac{[p + (1-p)t]^2 + (pt + 1-p)^2}{1+t} - \alpha \frac{1+t}{2} - (1-\alpha) \frac{2}{\pi} \mathcal{E} \left(\sqrt{1-t^2} \right) \right\} \\ &= a \left\{ \frac{(1-2p)^2 r^2 + 1}{1+r} - \alpha \frac{1}{1+r} - (1-\alpha) \frac{2}{\pi} \frac{2\mathcal{E} - (r')^2 \mathcal{K}}{1+r} \right\} \\ &= \frac{a}{1+r} \left[(1-2p)^2 r^2 + 1 - \alpha - (1-\alpha) \frac{2}{\pi} (2\mathcal{E} - (r')^2 \mathcal{K}) \right]. \end{aligned}$$

From this and Lemma 2.2, Theorem 3.1 follows. \square

Corollary 3.1. *For $r \in (0, 1)$ and $r' = \sqrt{1-r^2}$, we have*

$$\frac{\pi}{2} \left[\frac{17 + 30r' + 17(r')^2}{8(1+r')} - \frac{3(1+r')}{2} \right] < \mathcal{E}(r) < \pi \left[\frac{r' + 2(1-r')^2/\pi}{1+r'} \right]. \quad (3.2)$$

Proof. This follows from letting $\alpha = \frac{3}{4}$, $\lambda = \frac{5}{8}$, and $\mu = \frac{1}{2} \left(1 + \frac{\sqrt{4/\pi-1}}{2} \right)$ in Theorem 3.1. \square

4. REMARKS

Remark 4.1. Recently, the complete elliptic integrals have attracted attention of numerous mathematicians. In [9], it was established that

$$\begin{aligned} \frac{\pi}{2} \left[\frac{1}{2} \sqrt{\frac{1+(r')^2}{2}} + \frac{1+r'}{4} \right] &< \mathcal{E}(r) \\ &< \frac{\pi}{2} \left[\frac{4-\pi}{(\sqrt{2}-1)\pi} \sqrt{\frac{1+(r')^2}{2}} + \frac{(\sqrt{2}\pi-4)(1+r')}{2(\sqrt{2}-1)\pi} \right], \end{aligned} \quad (4.1)$$

for all $r \in (0, 1)$. In [11] it was proved that

$$\frac{\pi}{2} - \frac{1}{2} \log \frac{(1+r)^{1-r}}{(1-r)^{1+r}} < \mathcal{E}(r) < \frac{\pi-1}{2} + \frac{1-r^2}{4r} \log \frac{1+r}{1-r}, \quad (4.2)$$

for all $r \in (0, 1)$. In [18] it was presented that

$$\frac{\pi}{2} \frac{\sqrt{6+2\sqrt{1-r^2}-3r^2}}{2\sqrt{2}} \leq \mathcal{E}(r) \leq \frac{\pi}{2} \frac{\sqrt{10-2\sqrt{1-r^2}-5r^2}}{2\sqrt{2}} \quad (4.3)$$

for all $r \in (0, 1)$. In [9] it was pointed out that the bounds in (4.1) for $\mathcal{E}(r)$ are better than the bounds in (4.2) for some $r \in (0, 1)$.

Remark 4.2. The lower bound in (3.2) for $\mathcal{E}(r)$ is better than the lower bound in (4.1). Indeed,

$$\begin{aligned} \frac{17+30x+17x^2}{8(1+x)} - \frac{3(1+x)}{2} - \left[\frac{1}{2} \sqrt{\frac{1+x^2}{2}} + \frac{1+x}{4} \right] \\ = \frac{3x^2+2x+3-2\sqrt{2(1+x^2)}(1+x)}{8(1+x)} \end{aligned}$$

and

$$(3x^2+2x+3)^2 - [2\sqrt{2(1+x^2)}(1+x)]^2 = (1-x)^4 > 0$$

for all $x \in (0, 1)$.

Remark 4.3. The following equivalence relations show that the lower bound in (3.2) for $\mathcal{E}(r)$ is better than the lower bound in (4.3):

$$\begin{aligned} \frac{17+30x+17x^2}{8(1+x)} - \frac{3(1+x)}{2} &> \frac{\sqrt{6+2x-3(1-x^2)}}{2\sqrt{2}} \\ \iff (5x^2+6x+5)^2 &> 8(x+1)^2(3x^2+2x+3) \\ \iff (x-1)^4 &> 0, \end{aligned}$$

where $x \in (0, 1)$.

REFERENCES

- [1] H. Alzer and S.-L. Qiu, *Monotonicity theorems and inequalities for complete elliptic integrals*, J. Comput. Appl. Math. **172** (2004), no. 2, 289–312; Available online at <http://dx.doi.org/10.1016/j.cam.2004.02.009>. 2
- [2] G. D. Anderson, M.K. Vamanamurthy and M. Vuorinen, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, John Wiley & Sons, New York, 1997. 3
- [3] R. W. Barnard, K. Pearce, and K. C. Richards, *An inequality involving the generalized hypergeometric function and the arc length of an ellipse*, SIAM J. Math. Anal. **31** (2000), no. 3, 693–699; Available online at <http://dx.doi.org/10.1137/S0036141098341575>. 2

- [4] F. Bowman, *Introduction to Elliptic Functions with Applications*, 2nd ed., Dover Publications, New York, 1961. 2
- [5] P. F. Byrd and M. D. Friedman, *Handbook of Elliptic Integrals for Engineers and Scientists*, 2nd ed., Springer-Verlag, New York, 1971. 2
- [6] Y.-M. Chu and M.-K. Wang, *Inequalities between arithmetic-geometric, Gini, and Toader means*, Abstr. Appl. Anal. **2012**, Article ID 830585, 11 pages; Available online at <http://dx.doi.org/10.1155/2012/830585>. 2
- [7] Y.-M. Chu and M.-K. Wang, *Optimal Lehmer mean bounds for the Toader mean*, Results Math. **61** (2012), no. 3-4, 223–229; Available online at <http://dx.doi.org/10.1007/s00025-010-0090-9>. 2
- [8] Y.-M. Chu, M.-K. Wang, and X.-Y. Ma, *Sharp bounds for Toader mean in terms of contraharmonic mean with applications*, J. Math. Inequal. **7** (2013), no. 2, 161–166; Available online at <http://dx.doi.org/10.7153/jmi-07-15.pdf>. 2
- [9] Y.-M. Chu, M.-K. Wang, and S.-L. Qiu, *Optimal combinations bounds of root-square and arithmetic means for Toader mean*, Proc. Indian Acad. Sci. Math. Sci. **122** (2012), no. 1, 41–51; Available online at <http://dx.doi.org/10.1007/s12044-012-0062-y>. 2, 5
- [10] Y.-M. Chu, M.-K. Wang, S.-L. Qiu, and Y.-F. Qiu, *Sharp generalized Seiffert mean bounds for Toader mean*, Abstr. Appl. Anal. **2011** (2011), Article ID 605259, 8 pages; Available online at <http://dx.doi.org/10.1155/2011/605259>. 2
- [11] B.-N. Guo and F. Qi, *Some bounds for the complete elliptic integrals of the first and second kind*, Math. Inequal. Appl. **14** (2011), no. 2, 323–334; Available online at <http://dx.doi.org/10.7153/mia-14-26>. 2, 5
- [12] F. Qi, L.-H. Cui, and S.-L. Xu, *Some inequalities constructed by Tchebysheff's integral inequality*, Math. Inequal. Appl. **2** (1999), no. 4, 517–528; Available online at <http://dx.doi.org/10.7153/mia-02-42>. 2
- [13] F. Qi and Z. Huang, *Inequalities for complete elliptic integrals*, Tamkang J. Math. **29** (1998), no. 3, 165–169; Available online at <http://dx.doi.org/10.5556/j.tkjm.29.1998.165-169>. 2
- [14] F. Qi, D.-W. Niu, and B.-N. Guo, *Refinements, generalizations, and applications of Jordan's inequality and related problems*, J. Inequal. Appl. **2009** (2009), Article ID 271923, 52 pages; Available online at <http://dx.doi.org/10.1155/2009/271923>. 2
- [15] S.-L. Qiu and J.-M. Shen, *On two problems concerning means*, Journal of Hangzhou Institute of Electronic Engineering **17** (1997), no. 3, 1–7. (Chinese) 2
- [16] M. Vuorinen, *Hypergeometric functions in geometric function theory*, in: Special Functions and Differential Equations, Proceedings of a Workshop held at The Institute of Mathematical Sciences, Madras, India, January 13–24, 1997, Allied Publ., New Delhi, 1998, 119–126. 2
- [17] M.-K. Wang, Y.-M. Chu, S.-L. Qiu, and Y.-P. Jiang, *Bounds for the perimeter of an ellipse*, J. Approx. Theory **164** (2012), no. 7, 928–937; Available online at <http://dx.doi.org/10.1016/j.jat.2012.03.011>. 2
- [18] L. Yin and F. Qi, *Some inequalities for complete elliptic integrals*, available online at <http://arxiv.org/abs/1301.4385>. 5

(Wei-Dong Jiang) DEPARTMENT OF INFORMATION ENGINEERING, WEIHAI VOCATIONAL COLLEGE, WEIHAI CITY, SHANDONG PROVINCE, 264210, CHINA

E-mail address: jackjwd@hotmail.com

(Feng Qi) DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN CITY, 300387, CHINA

E-mail address: qifeng618@gmail.com, qifeng618@hotmail.com

URL: <http://qifeng618.wordpress.com>